# Pesticides in agricultural irrigation-return flow in the Columbia Basin Project, 2002-04

Lonna Frans
U.S. Geological Survey
September 21, 2006
Presented at Columbia River Toxics Reduction
Strategy Meeting




## **Outline**

- Study design
- > Summary of key results
  - Occurrence and distribution of pesticides
  - Timing of detections
  - Relation to other agricultural basins



- •4 sites
- •Samples collected from June 2002-October 2004
- 10 Samples during irrigation season and 2 during non-irrigation season
- Analyzed for nutrients, major ions, and 107 pesticides







#### Occurrence and distribution of pesticides

- •42 pesticides and 5 metabolites detected
- •Sand Hollow had the most compounds detected (37) and Crab Creek had the fewest compounds detected (28)
- Herbicides are more frequently detected than insecticides
- Concentrations of chlorpyrifos, lindane, azinphosmethyl and dinoseb exceeded aquatic-life criteria
















# Conclusions

- > 42 pesticides and 5 metabolites detected
- Sand Hollow had the most compounds detected (37) and Crab Creek had the least (28)
- Atrazine, bentazon, diuron and 2,4-D were most frequently detected herbicides
- Chlorpyrifos and azinphos-methyl were the most frequently detected insecticides



### Conclusions

- Total number of pesticides detected was higher during the irrigation season than during non-irrigation season
- 3 insecticides (azinphos-methyl, chlorpyrifos, and lindane) and 1 herbicide (dinoseb) were detected at concentrations that exceeded aquatic-life criteria
- Pesticides detected in this study were at lower concentrations but higher frequency than other agricultural basins around the Nation

